

## List of Figures

- Figure 1.1** Geological map of the northwestern Ladakh Himalaya (after Thakur and Rawat, 1992). The grey line shows the boundary between the Indus Molasse and Ladakh Batholith, red line with teeth show the faults and stars indicate the study sites.....5
- Figure 1.2** The structural cross-section along X-Y (shown in Fig 1.1), showing deformation of Indus basin sedimentary rocks (after Brookfield and Andrews-Speed, 1984). This small area (~35 km) has several north verging trusts and high angled reverse faults; Zanskar Back Thrust, Choksti thrust, and Upshi-Bazgo thrust.....14
- Figure 2.1** The energy level diagram explaining the TL/OSL working mechanism. In step-1, the ionization of the electrons occurred due to exposure of nuclear radiations. The formation of traps (T) and luminescence center (L) is due to movement of electrons and hole in the lattice of mineral, respectively. Step-2, represents the storage of electrons in the trap by continuous ionization. For dating, the life of trap should be more than the life of accumulation of electrons, therefore, the trap should not leak and hence the depth of trap should be equal to more than 1.6 eV. Step- 3, represents the eviction of electrons from the traps after get stimulated by suitable wavelength of light so that the electron get diffused into the conduction band and get attracted by the recombination center (luminescence center) to generate luminescence (Aitken, 1998).....25
- Figure. 2.2** Natural OSL and TL signals reduced in quartz and feldspar after exposure to sunlight. OSL has faster reduction than the TL in both quartz and feldspar (from Godfrey-Smith et al., 1988).....27

- Figure 2.3** Schematic representation of the building up and zeroing of the luminescence signals in a sediment sample. The erosional and transportation processes reduce the luminescence through direct exposure to sunlight or daylight. The build-up of luminescence is through accumulation of electrons in traps by getting flux of radiations due to continuous decay of U, Th and K and supply  $\alpha$ ,  $\beta$  and  $\gamma$  radiations. The cosmic radiations helps in building-up the luminescence signals in the sample (modified after Wintle and Huntley, 1982; Lepper et al., 2001).....28
- Figure 2.4** The schematic algorithm for SAR protocol.  $L_N$  and  $T_N$  are natural and test dose luminescence,  $L_1$  and  $T_1$  are luminescence measured after first incremental dose followed by test dose. This protocol is also known as five points SAR protocol because it has three incremental dose followed by one zero dose and one repetition of any incremental dose.....32
- Figure 2.5** The sensitivity corrected luminescence is plotted against the given incremental doses to generate growth curve. A straight line which cuts the growth curve from natural dose, when extended perpendicularly to dose axis, gives palaeodose. The inset shows the shine down curve for natural and test doses.....34
- Figure 2.6** The preheat plateau and dose recovery tests were subjected to LD-1047 to know the suitable preheat temperature for applicability of SAR protocol. (a) In preheat plateau test, the stability in the observed palaeodose at preheat temperatures between 200° and 240° C. At temperature lower than 180° C and higher than 240° C shows small gradient. This test suggest the temperature in between 200° - 240° C are suitable for SAR. (b) In dose recovery test, the given dose was extracted at different preheat temperature. At higher

temperatures, the palaeodose scattering and recuperation ratio exceeded from the acceptable limit. The lower temperatures having least recuperation ratio and less scattering, therefore 220° C is used as preheat temperature.....36

**Figure 2.7** (A) and (B) Histogram and weighted mean dose distribution showing partial bleaching of sample from Biamah section, (C) the rectangle in the radial plot representing the weighted mean palaeodoses used for age calculation, (D) the shine down curve, and (E) the growth curve for the same sample.....39

**Figure 2.8** Histogram and frequency distribution curve of sample from Saboo sand ramp are represented in (A) and (B). (C) In radial plot, the hollow rectangle bar represents the mean, whereas the shaded bar represents the least 20 % palaeodoses, which was considered for age calculation, (D) the shine down curve showing no feldspar contamination in this sample, and (E) the growth curve for the same sample.....40

**Figure 3.1** (A) Longitudinal river profile of the Indus River, showing studied sites. (B) Stream length gradient (SL) index variation with the downstream distance. The analysis of these two geomorphic tools identifies the whole reach into four segments.....49

**Figure 3.2** (A) Location map of Indus River. Small rectangle showing the Segment I. (B) Terrace map for Segment I. The yellow area shows the extension of fill terrace. (C) Valley cross section at Nyoma showing braided Indus River. (D) The google earth image showing the valley of the Indus at Nyoma is ~ 4 km wide. The closely spaced paleo-meander is also visible, implying increased channel gradient. (E) A panoramic view of the Indus at Nyoma, showing braided nature of the trunk channel.....55

- Figure 3.3** (A) Location map of Indus River and Segment-II. (B) A fill terrace in the segment. (C) The google earth image showing Indus Molasse as a bedrock at Mahe section. (D) Fill terraces, located on the right bank of the river as shown in valley cross-section.....57
- Figure 3.4** (A) Location map. (B) The goole earth showing the location of Kesar section along the Indus River. (C) A fill terrace has been marked in V-shaped valley. However, a debris flow has also been marked on the valley wall.....58
- Figure 3.5** (A) Location map. (B) A panoramic view of Niornis section showing the geomorphological setup. (C) A valley cross-section (A-A') showing a V-shaped valley and a fill terrace along the Indus River.....59
- Figure 3.6** (A) Location map. (B) At Kiari, the Indus River makes a sharp diversion in its path from the Indus Molasse to the batholith. The yellow line showing the contact of Indus Molasse and Ladakh Batholith. (C) Fill terrace in the V-shaped valley.....60
- Figure 3.7** (A) Location map of the Gaik section. (B) Google earth image showing, in a narrow deep gorge. A-A' is a valley cross-section at this section. (C) A valley fill terrace has been located in V-shaped valley.....62
- Figure 3.8** (A) Terrace map around Tirido section. (B) Google earth image, locating the studied section, valley cross-section (A-A') and Batholith as bedrock. (C) A valley cross-section A-A' indicates a V-shaped valley and a fill terrace on the left bank of the river. (D) A panoramic view of the Tirido section. The bedrock is grano-dioritic, where the river is flowing in a deep gorge.....63

- Figure 3.9** (A) Location and terrace map for Hymia sections. (B) Google earth image showing Hymia-1 and Hymia-2 sections. At these sections, the valley become slightly wider. The valley is mainly covered by large fans coming from both banks of the river. (C) and (D) Vally cross-sections for Hymia-1 and Hymia-2.....64
- Figure 3.10** (A) Regional map showing the Indus River and major localities in Ladakh. A box locating the Segment-III. (B) A terrace map of the Segment-III. The grey line is the molasse-batholith contact line, black line with teeth is Choksti thrust dipping south. (C) A topographic map of Segment-III, indicates the fans emerging from south are related to the activity along the Choksti thrust. The fans emerging from north, are belong to amphitheater valleys.....66
- Figure 3.11** (A) Terrace map locating the Upshi section. (B) The google earth image showing the topographic view around Upshi-R and Upshi-L sections. Here the Indus River is flowing on the contact of Indus Molasse and Ladakh Batholith, which is known as Upshi-Bazgo thrust. (C) The valley cross-section of both the sections indicating wider V-shaped valley, which is covered by alluvial fans.....68
- Figure 3.12** (A) Terrace map of Kharu section. (B) The google earth image shows that the river is flowing along the Upshi-Bazgo thrust. Alluvial fans emerges from the batholith and molasse side and unload the sediment into the trunk channel. (C) The valley cross-section of this section indicating that a wider V-shaped valley, where the topographic relief is low.....69
- Figure 3.13** (A) Terrace map. (B) A google earth image locating the Stakna, Stakna-1 and Stakna-2 sections and a topographic view around these sections. Stakna-1 section located just opposite bank on Stakna section. Stakna-2 is further downstream, where valley is covered by

alluvial fans from both sides of the river. (C) The valley cross-section at this section indicating the V-shaped valley become narrower due to alluvial fans.....71

**Figure 3.14** (A) Terrace map locating the Spituk section. (B) The google earth image shows a wide valley (~ 1200 m). The river follows a braided pattern. The banks of the trunk channel are not stable and bounded by fan in the south and bedrock in the north. (C) The valley cross-section at Spituk (Leh), indicates that the south bank of Indus is flanked by fanglomerate and north bank has Spituk paleolake deposits lying on fluvial sand.....72

**Figure 3.15** (A) A regional map for Segment-IV. (B) A panoramic view of this segment, where Indus is flowing in a deep gorge. (C) The terrace map of Segment-IV, which is divided into three parts; 1) Nimu to Saspol, 2) Saspol to Dumkhar, and 3) Dumkhar to Dah. The yellow, green and red fill along the Indus are filled, strath level – 1 and strath level – 2 respectively. The grey line shows the molasse-batholith contact.....74

**Figure 3.16** (A) Terrace map showing the location of the Nimu and the Saspol sections. (B) The google earth image locating the valley cross-section at Nimu, Upshi-Bazgo thrust and paleo-Indus courses. (C) and (D) The valley cross-section of Nimu and Saspol, indicates V-shaped valley with two levels of terraces. (E) The google earth image shows the Saspol section.....75

**Figure 3.17** (A) The terrace map of Segment-IV, from Saspol to Dumkhar, locating the fill (T-1) and strath (T-2) terraces in green and red colours, respectively. (B) The google image shows that the river flows in the molassic bedrock. The contact of molasse and batholith is shown by bold grey dotted line. At Nurla section, an arc shaped

depressed valley indicates the paleo-Indus course. (C) A panoramic view of the Indus gorge at the Nurla section, where T-2 is situated at 142 m arl. (D) A valley cross-section shows a deep V-shaped valley.....77

**Figure 3.18** (A) Terrace map locates Khalsi section. (B) The google earth image shows that the Indus is flowing over the molassic bedrock and formed a deep gorge. There are two level of strath terraces (T-1' and T-2) and one level of fill terrace (T-1). T-2 can be traceable on both banks of the Indus River at this section. (C) A panoramic view of T-2 terrace. (D) The valley cross-section shows T-2 is lying at ~ 179 m arl.....78

**Figure 3.19** (A) Terrace map locates Dumkhar section. (B) The google earth image shows that the Indus River follows a meandering pattern. At this section, the strath terrace (T-2) can be traceable downstream. (C) The V-shaped valley shows a deep gorge.....79

**Figure 3.20** (A) A terrace map along the Indus River, shows the studied sections Skyurbuchan Gompa and Skyurbuchan downstream. (B) In google earth image shows Skyurbuchan Gompa section, which lies in the molassic bedrock, whereas the Skyurbuchan downstream section lies in the batholith bedrock. (C) and (D) The valley cross-section of Skyurbuchan Gompa, and Skyurbuchan downstream sections respectively.....81

**Figure 3.21** (A) A terrace map shows the studied sections Biamah and Dah. (B) The google earth image shows straight channel of the Indus River, here the bedrock is Ladakh Batholith. (C) and (D) The valley cross-sections at Biamah and Dah, respectively. The strath level truncated near Biamah, because Indus entered into the batholith. The V-shaped valley with deep gorge indicating high erosion.....82

<b>Figure 3.22</b>	Location map shows 21 sand ramp sections in the Leh valley. Blue bold line represents the Indus River, where thin blue lines its tributaries, light grey are elevation contours. The ASTER DEM is used as the base map.....84
<b>Figure 3.23</b>	(A) A picture showing Spituk sand ramp section. (B) The ramp profile where bedrock is batholith; horizontal line with yellow base are aeolian sand; grey wavy beds are intra-dunal lacustrine deposits and horizontal parallel grey line are Spituk paleo-lake deposits....88
<b>Figure 3.24</b>	(A) A photograph showing the Saboo and ramp. (B) The Saboo ramp profile shows a composite deposits of aeolian, fluvial and talus environment.....89
<b>Figure 3.25</b>	(A) A photograph of Choglamsar sand ramp. (B) The ramp profile shows a composite deposits of aeolian, fluvial and talus sediment.....90
<b>Figure 3.26</b>	A photograph of Shey-2 sand ramp section. This section also has composite sediments of various depositional environment.....91
<b>Figure 3.27</b>	(A) A photograph of Shey-3 sand ramp section, shows talus, lacustrine and aeolian sediments. (B) The ramp profile, demarking the extension of different sedimentary environment in one composite section.....92
<b>Figure 4.1</b>	(A) Litholog showing clast supported horizontally stratified imbricated gravels (Gh) facies. The longer axis of discoid gravels are dipped to upstream direction, due to high energy of Indus River. (B) A photograph giving a view of the Gh facies at Stakna-1 section.....97



- Figure 4.2** (A) Litholog showing a thick succession of clast supported massive gravels (Gcm) facies. (B) The photograph showing accretion of this facies at Upshi-L section due to amalgamation of channel bars. ....98
- Figure 4.3** (A) The matrix supported graded gravels consist of sub-angular to poorly rounded, moderately to poorly sorted gravels. This facies is often capped by a sand facies. (B) Panoramic view of fan bound sequence showing the Gmg facies.....100
- Figure 4.4** (A) The well rounded, moderately to well sorted gravels of clast supported planar cross-bedded gravels (Gp). (B) The photograph showing Gp lithofacies exposed at Nurla section.....101
- Figure 4.5** (A) and (B) showing the litholog and photograph of horizontally laminated sand (Sh) lithofacies from Hymia section. The Sh facies is generally preceded by the gravel facies. (C) Showing a planar cross bedded sand facies (Sp), having sharp and erosional contacts with lower channel bound gravelly unit and upper fan bound unit, respectively at the Mahe section. (D) Photograph showing the same section showing Sp facies.....102
- Figure 4.6** Hymia-1 section having both channel bound and fan bound lithofacies associations. (A) Litholog of Hymia-1 section representing the association of Gh – Sh as a channel bound, and Gmg-Sh as a fan bound lithofacies association. (B) A picture of Hymia-1.....105
- Figure 4.7** A photograph of identified sand ramp facies: (A) Parallel laminated medium sand at Saboo, (B) Poorly sorted coarse sand, (C) Parallel bedded angular gravel, and (D) Parallel laminated clayey silt at Shey.....107

<b>Figure 4.8</b>	(A) Bivariate plot between mean size and standard deviation of Saboo, Kelso (Mojave Desert) and Ejina dunes (Chinese Desert) in Passega's curve (1964, 1977). (B) Comparison of mean size and standard deviation of these sand ramps. The narrow scattering characterizes the aeolian sand of Saboo and Leh valley. (C) The stereographic image of aeolian sand unit showing the angularity of quartz grains.....109
<b>Figure 4.9</b>	(A) Mean grain size, SD and clay % of Saboo sand ramp section (43 samples). (B) The environmental magnetic parameters of Saboo section. Top and middle crust showing sharp bulge in both grain size and environmental magnetic parameters. The shaded bars provide information regarding changes in depositional processes and climatic fluctuations.....111
<b>Figure 4.10</b>	SEM images: (A) un-weathered, perfect crystalline magnetite in the hard crust. (B) Weathered magnetite grain in the aeolian unit. ....112
<b>Figure 4.11</b>	Raman spectrum of top and middle hard crust along with parallel laminated sand unit showing presence of magnetite throughout the Saboo sand profile.....113
<b>Figure 4.12</b>	X-ray diffraction patterns of representative samples from top and middle hard crust along with parallel laminated sand unit showing dominance of illite and chlorite clay minerals throughout the Saboo sand profile.....114
<b>Figure 4.13</b>	(A) Both banks of the Indus River at Mahe showing valley fill in valley cross-section. The Mahe-L and Mahe-R, lithologs shown on the left and right banks of the valley cross-section, suggest that both channel and fan bound processes filled the valley. (B) A photograph of Mahe-R .....117

- Figure 4.14** (A) The valley cross-section of Kesar section. (B) A picture of fill terrace at Kesar showing, a sharp erosional contact between the lower and upper Gh facies.....119
- Figure 4.15** (A) Litholog and (B) valley cross-section showing the valley fill. (C) Photograph showing the channel bound facies association at Niornis.....120
- Figure 4.16** (A) The litholog and valley cross-section of the Kiari section. The litholog shows both fan and channel bound facies. (B) Photograph showing the OSL age from Sp facies.....121
- Figure 4.17** (A) Gaik section: litholog, valley cross-section (V-shaped). The litholog showing mixed and batholithic sourced clast. (B) and (C) suggest that the valley fill has been influenced by both channel and fan bound processes.....123
- Figure 4.18** (A) Litholog of Tirido section showing the aggradation of the narrow valley by both channel and fan bound activities in a narrow valley. (B) Photograph showing sedimentary fill in the Tirido....125
- Figure 4.19** (A) The valley at Hymia-1 is narrow and V-shaped, where the valley fill sequence lies on the left bank. The litholog showing mixed clasts and aggradation is followed by channel bound processes. (B) Photograph showing Hymia-1 section.....127
- Figure 4.20** (A) The valley cross-section and litholog suggest that fan bound process was very active at  $13\pm 2$  ka, where the channel was very active during  $7\pm 1$  ka. (B) Photograph showing channel bound facies in between the successive fan bound processes at Hymia-2, located near to second bridge (person for scale is encircled).....129

- Figure 4.21** (A) and (B) Represent the litholog and panoramic view of Upshi-R section. A 38 m Upshi-R section is a fill terrace made up of both channel and fan activities. (C) and (D) Showing channel and fan bound activities at Upshi-L section as shown in litholog and photograph. (E) and (F) The valley cross-sections of both Upshi-R and Upshi-L.....132
- Figure 4.22** (A) Litholog showing the 30 m thick section made up of channel bound processes, where the clasts are sourced from the fans coming from the Indus Molasse. The valley cross-section representing the width of the valley is mainly filled with fan sediment. (B) A photograph of Kharu.....134
- Figure 4.23** (A) A picture of Stakna. (B) Litholog and valley cross-section. (C) The litholog of Stakna-1 lies on right bank of the river just in front of the Stakna. (D) The google earth view of all three sections at Stakna. (E) The litholog of Stakna-2 section showing that fan bound processes were dominant at ~ 30 ka.....137
- Figure 4.24** (A) and (B) The litholog and valley cross-section of the Spituk lake section. (C) A picture showing the bottommost part of the Spituk section.....140
- Figure 4.25** (A) Cross-section showing the configuration of T-1 and T-2; the litholog of T-2 sequence showing a more prominent channel bound deposit at the Nimu gorge. (B) Panoramic view of Nimu section near the confluence of Indus and Zanskar rivers with fill and strath terraces T-1 and T-2, respectively.....141

- Figure 4.26** (A) The cross-section showing T-1 and T-2; the litholog shows channel bound deposit in both T-1 and T-2. (B) Panoramic view of the Saspol section.....144
- Figure 4.27** (A) A photograph of the Nurla section showing one fill (T-1) and one strath (T-2) terrace. (B) A closer view of T-1. (C) The lithologs of T-1 and T-2 showing that both terraces have aggraded via channel bound processes. The valley cross-section signifies the reduced valley width and deep and narrow gorge.....146
- Figure 4.28** (A) and (B) Showing closer view of strath terraces T-2 and T-1' indicating the channel bound LFA. (C) The lithologs of T-1, T-1' and T-2 representing aggradation via fluvial processes. The valley cross-section showing the arrangement of three levels of terraces in the wide valley.....148
- Figure 4.29** (A) The litholog of fill terrace T-1, showing the channel bound deposits T-2 is associated with both channel and fan bound deposits at Dumkhar. (B) and (C) Showing T-1 and T-2.....151
- Figure 4.30** (A) Lithologs of two terraces at the Skyurbuchan Gompa section suggest that the river aggraded via channel bound processes. (B) A photograph of T-2 terrace. (C) The valley cross-section showing wide valley at the time of T-2 aggradation and narrow at present day (T-1). (D) The litholog of T-1 and T-2 of the Skyurbuchan Downstream section. (E) A photograph of T-2 of the Skyurbuchan Downstream section. (F) V – shaped valley can easily be seen in the valley cross-section.....153
- Figure 4.31** Litholog of Biamah section. The valley is V – shaped and sediment lies on right bank of the river.....155

<b>Figure 4.32</b>	Litholog of Spituk sand ramp showing the facies distribution. ....	156
<b>Figure 4.33</b>	The litholog of Choglamsar sand ramp section showing the distribution of aeolian, fluvial and talus facies. ....	157
<b>Figure 4.34</b>	Parallel laminated medium sand, poorly sorted coarse sand and parallel bedded angular gravel facies that characterize the Saboo sand ramp. ....	160
<b>Figure 4.35</b>	The two lithologs of Shey-3 sand ramp showing the distribution of different facies. The lateral extension of these facies is quite similar as shown in (A) and (B). The litholog (A) is located ~ 50 m towards the river side from (B). This indicate that the intra-ramp-lake had limited lateral extension (~ 20 m). ....	162
<b>Figure 5.1</b>	(A) Clast supported gravels (Gh facies) showing imbrication angle ( $\theta$ ). (B) Photograph showing clast data: longest ( $D_L$ ), intermediate ( $D_I$ ) and shortest ( $D_S$ ) diameters, imbrication angle ( $\Theta$ ) and litho-type. ....	171
<b>Figure 5.2</b>	Inset map shows the location of sites and their valley cross-sections, where the paleo-discharge was estimated. ....	172
<b>Figure 5.3</b>	(A) Location map of Leh valley. (B) Google image shows Stakna section. From north of the Indus River, fans from batholith and from south, fans from Indus Molasse bound the valley. Here the valley width is measured 240 m. (C) Valley cross-section of Stakna section. (D) A photograph of Stakna section shows individual unit thickness, which represents the individual bar thickness and ultimately river depth (d). ....	173

<b>Figure 5.4</b>	A model showing the relation of intermediate clast size with its average velocity. Coefficient of correlation ( $R^2$ ) is ~ 80 %, which suggest strong correlation.....	175
<b>Figure 5.5</b>	Modern discharge of the Indus River from 2002 to 2006, at Leh station.....	179
<b>Figure 5.6</b>	A model showing Himalayan river response to glacial to interglacial climate change. The aggradation of a Himalayan rivers occurred during transient climate when sediment to water ratio was higher.....	183
<b>Figure 6.1</b>	The Indus River flowing SE to NW direction and draining various tectonic zones; the Karakoram fault zone in the north, the Indus Tsangpo Suture Zone, the northwestern Himalayan syntaxis and the major tributaries of Indus River.....	188
<b>Figure 6.2</b>	Aggradation in the Indus River vis-à-vis paleoclimate record. (A) Climate record from the Guliya Ice Core (after Thompson et al. 1997) and Benthic Stable Isotope Data from Arabian Sea sediment core NIOP905 87KYr (Jung et al., 2009). (B) Glacial record of Ladakh Himalaya, (C) OSL ages of the Channel and outwash fan sediments along the Indus River, (D) Frequency distribution curve of all the ages. Here it can be noted that the major aggradation has occurred during the warm MIS-1 and MIS-3.....	189
<b>Figure 6.3</b>	Major aggradation phases in the NE Himalayan to Nepal and the NW Himalayan Rivers.....	191
<b>Figure 6.4</b>	(A) Longitudinal river profile of Indus River, (B) SL index plotted along the Indus from upstream Nyoma to Dah Hanu section. (C) Concavity Index ( $K_s$ ) showing the variations (after Munack et al.,	

2014) in the SL index. A panel showing the lithology from the Indus River draining through its course. Based on longitudinal profile, SL and  $K_s$  indices, two major knickpoint have been observed (1) Mahe and (2) upstream Nimu section.....195

**Figure 6.5** (A) Longitudinal River profile showing valley fill and strath terraces in yellow and orange colours, respectively. (B) Longitudinal profile of Indus (Segment-IV) showing reconstructed two levels of paleo-river bed and ages of the strath terraces.....199

**Figure 6.6** Panel showing the bedrock incision rates in the Nanga Parbat Harmosh Massif (NPHM) region of the NW Himalayan Syntaxis (after Leland et al., 1998) and the incision rates deduced in this study. The distance zero represents the Indus – Zanskar confluence. Note the systematic downstream increase in the incision rates.....200

**Figure 6.7** Radial frequency plot of aspect of all 21 studied sand ramps in Leh valley.....205

**Figure 6.8** Shey-2, Shey-3, Choglamsar, Saboo and Spituk sections are correlated with major cold and warm events influenced by various workers in Ladakh and Thar Desert.....209